


# Air Mass RGB

## Quick Guide



### Why is the Air Mass RGB imagery important?

The Air Mass RGB is used to diagnose the environment surrounding synoptic systems by enhancing temperature and moisture characteristics of air masses. Cyclogenesis can be inferred by the identification of warm, dry, ozone-rich descending stratospheric air associated with jet streams and potential vorticity (PV) anomalies. The RGB can be used to validate the location of PV anomalies in model data. Additionally, this RGB can distinguish between polar and tropical air masses, especially along frontal boundaries and identify high-, mid-, and low-level clouds.



*Air Mass RGB from Himawari-8 AHI at 1000 UTC, 8 September 2015.*

### Air Mass RGB Recipe


Color	Band / Band Diff. (μm)	Physically Relates to...	<u>Small</u> contribution to pixel indicates...	<u>Large</u> contribution to pixel indicates...
Red	6.2 – 7.3	Vertical water vapor difference	Moist upper levels	<b>Dry upper levels</b>
Green	9.6 – 10.4	Tropopause height based on ozone	Low tropopause and high ozone	<b>High tropopause and low ozone</b>
Blue	6.2	Water vapor in ~200-500 mb layer	Dry upper levels or warm brightness temperatures	<b>Moist upper levels or cold brightness temperatures</b>

### Impact on Operations

#### Primary Application

#### **Identifying air masses:**

Polar and tropical air masses and the frontal boundary between them are readily seen in the RGB imagery.




**Inferring cyclogenesis:** It is relatively easy to see jet streams and stratospheric air intrusions with high PV, and the cyclonic activity created by these dynamics.

#### **Secondary Applications:**

Moisture boundaries, distinguishing between warm air masses with high and relatively low moisture, distinguishing between high clouds and mid-level clouds

### Limitations

**Limb effects:** The use of longer wavelength channels results in more atmospheric absorption at large viewing angles. As a result of the greater absorption, cooler brightness temperatures are measured. This limb cooling causes false blue and violet colors along the entire limb.



**Upper troposphere only:** Conditions in the mid- to upper troposphere can be detected but surface conditions cannot be directly observed.

**Cloud cover:** High cloud cover can obscure air masses and make interpretation more difficult.



# Air Mass RGB


## Quick Guide



### RGB Interpretation

- 1** Jet stream / PV / deformation zones / dry upper level (dark red/orange)
- 2** Cold air mass (dark blue/purple)
- 3** Warm air mass (green)
- 4** Warm air mass, less moisture (dull green)
- 5** High thick cloud (white)
- 6** Mid-level cloud (tan/salmon)
- 7** Low-level cloud (green, dark blue)
- 8** Limb effects (purple/blue)

*Note: colors may vary diurnally, seasonally, and latitudinally*




Air Mass RGB from Himawari-8 AHI at 1200 UTC, 15 November 2015.

### Usage with other products:

The 6.9  $\mu\text{m}$  water vapor channel (right) can be applied to observe air mass interactions, jet streaks, and deformation zones. In this image, from the same time as the RGB above, the PV anomaly is apparent, but air mass temps and moisture are not.

### AHI 6.9 $\mu\text{m}$ water vapor (At same time as above)



### Resources

UCAR/COMET  
[Multispectral Satellite Applications: RGB Products Explained.](#)

NASA/SPoRT  
[Aviation Forecasting RGB Products](#)

EUMETrain  
[RGB Interpretation Guide](#)